arXiv:1503.01141v1 [math.GM] 3 Mar 2015

On the Complete Evaluation of Jacobi Theta Functions

N.D.Bagis
Stenimahou 5 Edessa
Pellas 58200, Greece
email-nikosbagis@hotmail.gr

Abstract

Using numerical methods we construct evaluation formulas for the Jacobi θ_4 functions. Our results are conjectures, but are verified numerically.

1 Introduction

Let $K(x)$ be the complete elliptic integral of the first kind

$$K(x) = \frac{\pi}{2} {}_2F_1\left(\frac{1}{2}, \frac{1}{2}; 1; x^2\right) \tag{1}$$

and k_r, the elliptic singular modulus, solution of the equation

$$\frac{K\left(\sqrt{1-k_r^2}\right)}{K(k_r)} = \sqrt{r} \tag{2}$$

When r is positive rational the function k_r take algebraic values.
The 4th-Jacobian theta function is given by

$$\vartheta_4(z, q) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2} e^{2niz} \tag{3}$$

where $|q| < 1$ and $z \in \mathbf{C}$. Avoiding this definition we will use the equivalent notation

$$\theta(a, b; q) := \sum_{n=-\infty}^{\infty} (-1)^n q^{an^2+bn} \tag{4}$$

and we shall restrict to the case of a positive rational and b general rational.
In [Bag,Glas] we have shown that if $|q| < 1$ and

$$A(a, p; q) := q^{\frac{p}{12}-\frac{a}{2}+\frac{a^2}{2p}} \prod_{n=1}^{\infty} \left(1 - q^{np+a}\right)\left(1 - q^{np+p-a}\right) \tag{5}$$

with $p > 0$, then

$$A(a, p; q) = \frac{q^{\frac{p}{12}-\frac{a}{2}+\frac{a^2}{2p}}}{\eta(q^p)} \theta\left(\frac{p}{2}, \frac{p}{2} - a; q\right) \tag{6}$$

where η is the Dedekind-Ramanujan eta function

$$\eta(q) := \prod_{n=1}^{\infty}(1-q^n),\ |q|<1 \tag{7}$$

The above equality (6) is simple consequence of the Jacobi triple product formula (see [2]).

2 Conjecture

Having define what we need we state our

Conjecture.
Let $q = e^{-\pi\sqrt{r}}$ with $r > 0$, then for a, p rationals and $p > 0$ there always exist algebraic function $Q(x)$, $0 < x < 1$, such that

$$A(a,p,q) = Q(k_r),\ \forall r > 0 \tag{8}$$

Corollary.
If $q = e^{-\pi\sqrt{r}}$ and $r, |a|, p \in \mathbf{Q}_+^*$, then

$$A(a,p;q) = Algebraic\ Number \tag{9}$$

Some verifications (proofs) of the above conjecture have given in [8] for the case of theta functions of the form

$$\sum_{n=-\infty}^{\infty} q^{n^2+mn},\ m \in \mathbf{Z} \tag{10}$$

Moreover it has been shown that:
i) if $m = 2s$ (even), then

$$\sum_{n=-\infty}^{\infty} q^{n^2+2sn} = q^{-s^2}\sqrt{\frac{2K(k_r)}{\pi}} \tag{11}$$

ii) if $m = 2s+1$ (odd), then

$$\sum_{n=-\infty}^{\infty} q^{n^2+(2s+1)n} = 2^{5/6}q^{-(2s+1)^2/4}\frac{(k_{11}k_{12}k_{21})^{1/6}}{k_{22}^{1/3}}\sqrt{\frac{K(k_{11})}{\pi}} \tag{12}$$

where $k_{11} = k_r$, $k_{12} = \sqrt{1-k_{11}^2}$, $k_{21} = \frac{2-k_{11}^2-2k_{12}}{k_{11}^2}$, $k_{22} = \sqrt{1-k_{21}^2}$ and in view of the evaluation formula (see [8]):

$$\eta(q)^8 = \frac{2^{8/3}}{\pi^4}q^{-1/3}k_r^{2/3}(k_r')^{8/3}K(k_r)^4 \tag{13}$$

Examples of the above conjecture can also be found if we consider the function

$$r = k_i(x) := \left(\frac{K\left(\sqrt{1-x^2}\right)}{K(x)}\right)^2 \tag{14}$$

which, is the inverse function of the singular modulus k_r. Our method consist of inserting the value $r = k_i\left(\frac{m}{n}\right)$, where $0 < m < n$, m, n integers into the form $A(a, p; q)$ and get numerically, using the routine *RootApproximant* of the program Mathematica, a minimal polynomial which is esentialy the value of an algebraic number. This lead us to conclude that beneath (any theta function) exists minimal polynomials with coefficients rational polynomials of k_r (in all cases if Conjecture holds) and for fixed a, p, unique algebraic function $Q_{\{a,p\}}(x)$, such that these polynomials have solution the functions $Q_{\{a,p\}}(k_r)$.
A very easy example to see someone is with $a = 1$ and $p = 4$. In this case all the values of $A\left(1, 4, e^{-\pi\sqrt{k_i\left(\frac{m}{n}\right)}}\right)^{24}$ are rationals. With a simple algorithm one can see that

$$A\left(1, 4, e^{-\pi\sqrt{k_i(r)}}\right)^{24} = \frac{16(1-r^2)}{r^2} \tag{15}$$

Hence

$$A\left(1, 4, e^{-\pi\sqrt{r}}\right) = \sqrt[24]{\frac{16(1-k_r^2)}{k_r^2}} \tag{16}$$

and by relation (6)

Theorem 1.
If $q = e^{-\pi\sqrt{r}}$, $r > 0$

$$\sum_{n=-\infty}^{\infty}(-1)^n q^{2n^2+n} = q^{1/24}\eta(q^4)\sqrt[12]{\frac{4(1-k_r^2)}{k_r}}, \forall r > 0 \tag{17}$$

The continuation follows from the validity of (6) and (8) in rationals which are dense in reals.
Another example is with $p = 2$ and $a = 1/2$, where we have

$$A\left(1/2, 2, e^{-\pi\sqrt{r}}\right) = \sqrt[24]{\frac{4(1-k_r)^4}{k_r(1+k_r)^2}} \tag{18}$$

and is the same theta function as (16) by changing $q \to q^{1/2}$. For avoiding these cases it is useful to know that

$$\sum_{n=-\infty}^{\infty}(-1)^n q^{an^2+bn} = \sum_{n=-\infty}^{\infty} q^{an^2-bn} \tag{19}$$

and also that if $s > 0$

$$\sum_{n=-\infty}^{\infty}(-1)^n q^{an^2+bn}, \quad \sum_{n=-\infty}^{\infty}(-1)^n q^{asn^2+bsn} \tag{20}$$

are equivalent.

Theorem 2.
For $q = e^{-\pi\sqrt{r}}$, $r > 0$, we have

$$\sum_{n=-\infty}^{\infty} (-1)^n q^{2n^2+3n/2} = q^{-11/96}\eta(q^4)\sqrt[48]{\frac{4(1-k_r)^4(2+k_r-2\sqrt{1+k_r})^{12}}{k_r^{13}(1+k_r)^2}} \tag{21}$$

3 The Algorithm

In this section we give the algorithm for finding the expression $Q_{\{a,p\}}\left(k_r^2\right)$. Our method is based on interpolation. We find the minimal polynomial

$$P\left(A(a,p;q),m(q)\right) = 0, \tag{22}$$

with $m(q) = k_r^2$ and then solve with respect to $A(a,p;m(q))$, if $P(x,y) = 0$ is solvable.

The algorithm (In Mathematica Program)

$$Clear[A]$$

$$eta[q] := QPochhammer[q,q]$$

$$A[a,p,q] := q^{p/12-a/2+a^2/(2p)}eta[q^p]^{-1}Sum[(-1)^n q^{n^2p/2+(p-2a)n/2},\{n,-100,100\}]$$

$$Clear[q,x,y,u,v]$$

$$m[q] := InverseEllipticNomeQ[q]$$

$$x = Series[A[a,p,q]^{12},\{q,0,M\}];$$

$$y = Series[m[q],\{q,0,M\}];$$

$$t = Table[Coefficient[Sum[c[i,j]x^iy^j,q^n] == 0,\{n,1,s^2\}];$$

$$rr = Table[c[i,j],\{i,0,s\},\{j,0,s\}];$$

$$rr1 = Table[u^iv^j,\{i,0,s\},\{j,0,s\}];$$

$$mm = Normal[Extract[CoefficientArrays[t//Flatten, rr//Flatten],2]];$$

$$m0 = Normal[mm];$$

$$r1 = Take[NullSpace[m0],1].Flatten[rr]$$

$$Take[NullSpace[m0],1].Flatten[rr1]//Factor$$

4 Theoretical Results and Directions

In this section we will try to characterize these functions $Q_{\{a,p\}}(x)$. For this, assume that P_n is the n-th modular equation of $A(a,p;q)$, then

$$A(a,p;q^n) = P_n\left(A(a,p;q)\right) \tag{23}$$

Also assume that our conjecture (8) hold, then

$$Q_{\{a,p\}}\left(k_{n^2 r}\right) = P_n\left(Q_{\{a,p\}}(k_r)\right)$$

By using (14), we get

$$Q_{\{a,p\}}\left(k_{n^2 k_i(x)}\right) = P_n\left(Q_{\{a,p\}}(x)\right)$$

Setting

$$S_n(x) := k_{n^2 k_i(x)} \tag{24}$$

we have the next

Theorem 3
If the n-th modular equation of $A(a,p;q)$ is that of (23), then

$$k_{n^2 k_i(x)} = S_n(x) = {Q_{\{a,p\}}}^{(-1)}\left(P_n\left(Q_{\{a,p\}}(x)\right)\right),\ n = 2,3,4,... \tag{25}$$

If one manages to solve equation (25) with respect to $Q_{\{a,p\}}(x)$ for given a,p, then

$$\sum_{n=-\infty}^{\infty} (-1)^n q^{pn^2/2+(p-2a)n/2} = q^{-\frac{p}{12}+\frac{a}{2}-\frac{a^2}{2p}} \eta(q^p) Q_{\{a,p\}}(k_r), \forall r > 0 \tag{26}$$

and $Q_{\{a,p\}}(x)$ will be a root of a minimal polynomial of degree $\nu = \nu(a,p,x)$. Note that in case of rational $x \in (0,1)$ and a,p rational with $0 < a,p$, then the degree ν is independent of x and the minimal polynomial of $Q_{\{a,p\}}(x)$ will have integer coefficients.

Example.
The 2nd degree modular equation of $A(1,4;q)$ is

$$16u^8 + u^{16}v^8 - v^{16} = 0 \tag{27}$$

If we solve with respect to v we get $v = P_2(u)$, where $v = A(1,4;q^2)$ and $u = A(1,4;q)$. Moreover

$$P_2(w) = \frac{\left(w^{16} + w^4\sqrt{64 + w^{24}}\right)^{1/8}}{2^{1/8}} \tag{28}$$

It is $n=2$ and hold (see [9])

$$k_{4r}=\frac{1-\sqrt{1-k_r^2}}{1+\sqrt{1-k_r^2}} \tag{29}$$

Hence

$$S_2(x)=k_{4k_i(x)}=\frac{1-\sqrt{1-x^2}}{1+\sqrt{1-x^2}} \tag{30}$$

Finally we get from the relation (25) of Theorem 4:

$$Q^{(-1)}_{\{1,4\}}\left(\frac{\sqrt[8]{Q_{\{1,4\}}(x)^{16}+Q_{\{1,4\}}(x)^4\sqrt{Q_{\{1,4\}}(x)^{24}+64}}}{\sqrt[8]{2}}\right)=\frac{1-\sqrt{1-x^2}}{1+\sqrt{1-x^2}} \tag{31}$$

which has indeed a solution

$$Q_{\{1,4\}}(x)=\sqrt[12]{\frac{4(1-x^2)}{x}}.$$

Note.
We note that function $m(q)=k_r^2$ is implemented in program Mathematica. However a useful expansion is

$$k_r=\sqrt{m(q)}=4q^{1/2}\exp\left(-4\sum_{n=1}^{\infty}q^n\sum_{d|n}\frac{(-1)^{d+n/d}}{d}\right), \tag{32}$$

where $q=e^{-\pi\sqrt{r}}$, $r>0$.

5 Table

Here we give a table of evaluations, which does not include Theorems 1 and 2, for certain lower values a,p of theta functions.

1.

$$\sum_{n=-\infty}^{\infty}(-1)^n q^{3/2n^2+n/2}=q^{1/12}\eta(q^3)Q(k_r^2) \tag{33}$$

The polynomial which relates $u=A(1,3,q)^{12}$ with $v=m(q)=k_r^2$ is

$$u^4v^5-4u^4v^4+6u^4v^3-4u^4v^2+u^4v-16u^3v^6+84u^3v^5-12480u^3v^4-$$

$$-40712u^3v^3-12480u^3v^2+84u^3v-16u^3+196830u^2v^5-787320u^2v^4+1180980u^2v^3-$$

$$-787320u^2v^2+196830u^2v+19131876uv^5-76527504uv^4+114791256uv^3-$$

$$-76527504uv^2+19131876uv+387420489v^5-1549681956v^4+2324522934v^3-$$

$$-1549681956v^2 + 387420489v = 0 \tag{34}$$

2.

$$\sum_{n=-\infty}^{\infty} (-1)^n q^{3n^2-5n} = q^{-11/6}\eta(q^6)Q(k_r) \tag{35}$$

The polynomial equation which relates $u = A(8,6;q)^6$ with $v = m(q)^{1/2} = k_r$ is

$$u^8v^4 - u^8v^2 + 16u^6v^6 - 24u^6v^4 - 24u^6v^2 + 16u^6 - 486u^4v^4 +$$

$$+486u^4v^2 - 19683v^4 + 19683v^2 = 0 \tag{36}$$

3.

$$\sum_{n=-\infty}^{\infty} (-1)^n q^{3n^2+4n} = q^{-13/12}\eta(q^6)Q(k_r) \tag{37}$$

The polynomial equation which relates $u = A(-1,6,q)^6$, with $v = m(q)^{1/2} = k_r$ is

$$u^4v^3 - u^4v + 16u^3v^2 - 18u^2v^3 + 18u^2v + 4uv^4 - 8uv^2 + 4u + v^3 - v = 0 \tag{38}$$

4.

$$\sum_{n=-\infty}^{\infty} (-1)^n q^{4n^2+6n} = q^{-23/12}\eta(q^8)Q(k_r) \tag{39}$$

The polynomial equation which relates $u = A(-2,8;q)^{12}$, with $v = m(q^2)^2 = k_{4r}^4$ is

$$-u^4v - 64u^2v + 256v^2 - 512v + 256 = 0 \tag{40}$$

5.

$$\sum_{n=-\infty}^{\infty} (-1)^n q^{5/2n^2+3/2n} = q^{-1/60}\eta(q^5)Q(\eta_5(q^4)^5) \tag{41}$$

where

$$\eta_5(q) = \frac{1}{2}\left(-1 - h_5(q) + \sqrt{5 + 2h_5(q) + h_5(q)^2}\right) \tag{42}$$

and

$$h_5(q) = \frac{\eta(q^{1/5})}{q^{1/5}\eta(q^5)} \tag{43}$$

The polynomial equation which relates $u = A(1,5,q^2)^{15}$ with $v = \eta_5(q^4)^5$ is

$$u^4 + v^{11} + 55v^{10} + 1205v^9 + 13090v^8 + 69585v^7 + 134761v^6 - 69585v^5 +$$

$$+13090v^4 - 1205v^3 + 55v^2 - v = 0 \tag{44}$$

Note that

$$h_5(q^5)=\frac{\eta(q)}{q\eta(q^{25})}=\frac{1}{\sqrt{M_5(q)M_5(q^5)}}\left(\frac{m(q)}{m(q^{25})}\right)^{1/24}\left(\frac{m'(q)}{m'(q^{25})}\right)^{1/6}$$

is function of $k_r=\sqrt{m(q)}$, since (see [14])

$$\left(5M_5(q)-1\right)^5\left(1-M_5(q)\right)=256\cdot m(q)m'(q)M_5(q) \tag{45}$$

and $m'(q)=1-m(q)$.

For tables of singular modulus one can see [10],[11],[14].

References

[1]: M.Abramowitz and I.A.Stegun. 'Handbook of Mathematical Functions'. 9th ed., New York: Dover Pub. (1972)

[2]: J.V. Armitage and W.F. Eberlein. 'Elliptic Functions'. 2006 ed., New York: Cambridge University, (2006)

[3]: Nikos Bagis. 'On a general sextic equation solved by the Rogers-Ramanujan continued fraction'. arXiv:1111.6023v2, (2012)

[4]: Nikos Bagis. 'Evaluation of Fifth Degree Elliptic Singular Moduli'. arXiv:1202.6246v3, (2015)

[5]: N.D. Bagis and M.L. Glasser. 'Ramanujan type $1/\pi$ Approximation Formulas'. Journal of Number Theory, Vol(133), 2013 pp. 3453-3469.

[6]: Nikos Bagis. 'Parametric Evaluations of the Rogers Ramanujan Continued Fraction'. International Journal of Mathematics and Mathematical Sciences. Vol(2011). doi:101155/2011/940839.

[7]: Nikos Bagis and M.L. Glasser. 'Conjectures on the evaluation of alternative modular bases and formulas approximating $1/\pi$'. Journal of Number Theory, Vol(132), 2012 pp. 2353-2370.

[8]: Nikos Bagis and M.L. Glasser. 'Jacobian Elliptic Functions, Continued Fractions and Ramanujan Quantities'. arXiv:1001.2660v1, (2010)

[9]: B.C. Berndt. 'Ramanujan's Notedbooks Part III'. 1991 ed., New York: Springer-Verlag. (1991)

[10]: B.C. Berndt. 'Ramanujan's Notedbooks Part V'. 1998 ed., New York: Springer-Verlag. (1998)

[11]: J.M. Borwein and P.B. Borwein. 'Pi and the AGM'. 1987 ed., New York: John Wiley and Sons Inc., (1987)

[12]: D. Broadhurst. 'Solutions by Radicals at Singular Values k_N from New Class Invariants for $N \equiv 3 mod 8$'. arXiv:0807.2976 (math-phy), (2008)

[13]: ET. Whittaker and G.N. Watson. 'A course on Modern Analysis'. 4th ed., Cambridge University Press. (1927)

[14]: J.M. Borwein, M.L. Glasser, R.C. McPhedran, J.G. Wan, I.J. Zucker. 'Lattice Sums Then and Now'. Cambridge University Press. New York, (2013).

[15]: N.D. Bagis. 'On Generalized Integrals and Ramanujan-Jacobi Special Functions'. arXiv:1309.7247v2